


## Vectors

A vector is a mathematical object consisting of a magnitude (size) and a direction.


A vector can be represented graphically by an arrow:



A vector quantity is written in bold ( $\mathbf{A}$ ) or with a little arrow overhead ( $\vec{A}$ )


$A$  (no arrow, not bold) =  $|\vec{A}|$  = magnitude of the vector = positive number (magnitudes are positive by definition)

Examples of vector quantities: position, velocity, acceleration, force, electric field.



If two vectors have the same direction and the same magnitude, then they are the same vector



Vector = magnitude + direction (not location)



In 2D, we need 2 numbers to specify a vector  $\vec{A}$  :

- magnitude  $A$  and angle  $\theta$   
or
- components  $A_x$  and  $A_y$  (more on components later)


## Addition of Vectors




$$\vec{A} + \vec{B} = \vec{C}$$

Vector addition is commutative:  $\vec{A} + \vec{B} = \vec{B} + \vec{A}$


Graphical addition: "tip-to-tail" or "tail-to-head" method:



Addition by "parallelogram method" (same result as tip-to-tail method)




Can add lots of vectors (like steps in a treasure map: "take 20 steps east, then 15 steps northwest, then...")





$$\vec{A} + \vec{B} + \vec{C} + \vec{D} + \vec{E} = \vec{S}$$

Definition of negative of vector (same size, opposite direction):



Definition of multiplication of a vector by a number:



What about multiplication of a vector by a vector?


There are two different ways to define multiplication of two vectors:

(1) Dot product or scalar product  $\vec{A} \cdot \vec{B}$  and (2) Cross product  $\vec{A} \times \vec{B}$   
These will be defined later.


**Vector subtraction:**

$\vec{A} - \vec{B} = \vec{A} + (-\vec{B})$  ← "subtract" means "add negative of"

Graphically:  $\vec{D} = \vec{A} - \vec{B}$



**Components of a Vector**




$\vec{A} = A_x \hat{x} + A_y \hat{y}$

( $\hat{x}$  = "x-hat" is the *unit vector*, explained below)


$A_x = A \cos \theta$  = x-component = "projection of A onto x-axis"

$A_y = A \sin \theta$  = y-component = "projection of A onto y-axis"

Think of the  $A_x$  as the "shadow" or "projection" of the vector **A** cast onto the x-axis by a distant light source directly "overhead" in the direction of +y.




Components are numbers, not vectors. They do not have a direction, but they do have a *sign*, a (+) or (-) sign. If the "shadow" onto the x-axis points in the +x direction, then  $A_x$  is positive.



Here,  $B_x$  is negative, because the x-projection is along the -x direction.

$B_y$  is positive, because the y-projection is along the +y direction.



$$\cos\theta = \frac{A_x}{A} \Rightarrow A_x = A \cos\theta$$


$$\sin\theta = \frac{A_y}{A} \Rightarrow A_y = A \sin\theta$$

$$A = \sqrt{A_x^2 + A_y^2} \qquad \tan\theta = \frac{A_y}{A_x}$$

Magnitude  $A = |\vec{A}|$  is positive always, but  $A_x$  and  $A_y$  can be + or - .

**Unit Vectors**


A unit vector is a vector with magnitude = 1 (unity). Notation: a unit vector is always written with a caret (^) on top. The unit vectors  $\hat{x}$ ,  $\hat{y}$ , and  $\hat{z}$ , also written  $\hat{i}$ ,  $\hat{j}$ , and  $\hat{k}$ , are the unit vectors that point along the positive x-direction, y-direction and z-direction, respectively.




Any vector can be written in terms of its components like so:

$$\vec{A} = A_x \hat{i} + A_y \hat{j}$$

For instance, if  $A_x = 2$ ,  $A_y = -3$ , then the vector looks like :



**Example of vector math:**  $\vec{A} = 2\hat{i} - 3\hat{j}$ , meaning  $A_x = +2$ ,  $A_y = -3$  What is the magnitude A, and the angle  $\alpha$  of the vector with the positive x-direction ?



$$A = \sqrt{A_x^2 + A_y^2} = \sqrt{2^2 + 3^2} = \sqrt{4+9} = \sqrt{13} \simeq 3.6$$

$$\tan\alpha = \frac{3}{2} \Rightarrow \alpha = \tan^{-1}\left(\frac{3}{2}\right) = 56.3^\circ$$


**Vector Addition by Components:**

$$\vec{C} = \vec{A} + \vec{B} \Rightarrow$$

$$C_x = A_x + B_x$$

$$C_y = A_y + B_y$$

Proof by diagram:



Similarly, **subtraction by components:**


$$\vec{D} = \vec{A} - \vec{B} \Rightarrow D_x = A_x - B_x$$

$$D_y = A_y - B_y$$

**Position, Velocity, and Acceleration Vectors**

Velocity is a vector quantity; it has a magnitude, called the speed, and a direction, which is the direction of motion. Position is also a vector quantity. Huh? What do we mean by the magnitude and direction of position? How can position have a direction?

In order to specify the position of something, we must give its location in some coordinate system, that is, its location relative to some origin. We define the position vector  $\mathbf{r}$  as the vector which stretches from the origin of our coordinate system to the location of the object. The x- and y-components of the position vector are simply the x and y coordinates of the position. Notice that the position vector depends on the coordinate system that we have chosen.






If the object is moving, the position vector is a function of time  $\mathbf{r} = \mathbf{r}(t)$ . Consider the position vector at two different times  $t_1$  and  $t_2$ , separated by a short time interval  $\Delta t = t_2 - t_1$ . ( $\Delta t$  is read "delta-t") The position vector is initially  $\mathbf{r}_1$ , and a short time later it is  $\mathbf{r}_2$ . The change in position during the interval  $\Delta t$  is the vector  $\Delta \mathbf{r} = \mathbf{r}_2 - \mathbf{r}_1$ . Notice that, although  $\mathbf{r}_1$  and  $\mathbf{r}_2$  depend

on the choice of the origin, the change in position  $\Delta \mathbf{r} = \mathbf{r}_2 - \mathbf{r}_1$  is independent of choice of origin. Also, notice that change in something = final something – initial something.

In 2D or 3D, we define the velocity vector as  $\vec{v} = \lim_{\Delta t \rightarrow 0} \frac{\Delta \vec{r}}{\Delta t}$ .

As  $\Delta t$  gets smaller and smaller,  $\mathbf{r}_2$  is getting closer and closer to  $\mathbf{r}_1$ , and  $\Delta \mathbf{r}$  is becoming tangent to the path of the object. Note that the velocity  $\mathbf{v}$  is in the same direction as the infinitesimal  $\Delta \mathbf{r}$ , since the vector  $\mathbf{v}$  is a positive number ( $1/\Delta t$ ) times the vector  $\Delta \mathbf{r}$ . Therefore, the velocity vector, like the infinitesimal  $\Delta \mathbf{r}$ , is always tangent to the trajectory of the object.



The vector equation  $\vec{v} = \lim_{\Delta t \rightarrow 0} \frac{\Delta \vec{r}}{\Delta t}$  has x- and y-components. The component equations are

$$v_x = \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t}, \quad v_y = \lim_{\Delta t \rightarrow 0} \frac{\Delta y}{\Delta t}.$$

Any vector equation, like  $\vec{A} = \vec{B} + \vec{C}$ , is short-hand notation for 2 or 3 component equations:  $A_x = B_x + C_x, A_y = B_y + C_y, A_z = B_z + C_z$

The change in velocity between two times  $t_1$  and  $t_2$  is  $\Delta \mathbf{v} = \mathbf{v}_2 - \mathbf{v}_1$  (remember that change is

always final minus initial). We define the acceleration vector as  $\vec{a} = \lim_{\Delta t \rightarrow 0} \frac{\Delta \vec{v}}{\Delta t}$ . As we

mentioned in the chapter on 1D motion, the direction of the acceleration is the same as the direction of  $\Delta \mathbf{v}$ . The direction of the acceleration is NOT the direction of the velocity, it is the “direction towards which the velocity is tending”, that is, the direction of  $\Delta \mathbf{v}$ .

We will get more experience thinking about the velocity and acceleration vectors in the next few chapters.