


Putting Local Realism to the Test



“We can't solve problems by using the same kind of thinking we used when we created them.”

- Albert Einstein

Day 39:

Questions?

Revisit EPR-Argument

Testing Local Realism

Single Photon

Up Next:

Readings!

Finish Single-Photon Experiments


Wave-Particle Duality

Recently:

1. Hidden variables, locality, quantum interpretations.
2. Entanglement

Today:

1. Revisit the EPR argument.
2. Testing local realism
3. Single photon




What would be the expectation (average) value for m_x ?

$$\begin{aligned}\langle m_x \rangle &= P[|\uparrow_x\rangle](+m_B) + P[|\downarrow_x\rangle](-m_B) \\ &= (0.50)(+m_B) + (0.50)(-m_B) = 0\end{aligned}$$

Which interpretation sounds most reasonable to you?

- A) **Interpretation One:** An atom with a definite value of m_z also has a definite value of m_x , but that value changes so rapidly that we can't predict it ahead of time.
- B) **Interpretation Two:** An atom with a definite value of m_z also has a definite value of m_x but measuring m_z disturbs the value of m_x in some unpredictable way.
- C) **Interpretation Three:** An atom with a definite value of m_z doesn't have a definite value of m_x until measured.
- D) **A & B seem equally reasonable.**
- E) **Something else...**


Experiment One



The diagram shows a central box labeled 'source' with two arrows pointing outwards to the left and right. On the left is a box labeled '1' with a '+' sign in the top-left corner, a '-' sign in the bottom-left corner, and an upward-pointing arrow in the center. On the right is a box labeled '2' with a '-' sign in the bottom-right corner, a '+' sign in the top-right corner, and a downward-pointing arrow in the center.

- The results of **Experiment One** show that the measurements performed on **Atom 1** and on **Atom 2** are **anti-correlated**.
- **Anti**-correlated means that, whatever answer we get for **Atom 1**, we'll get the **opposite** answer for **Atom 2**, as long as we're asking the **same question**.
- Atom pairs in a correlated state $|\Psi_{12}\rangle = |\uparrow_1\rangle|\downarrow_2\rangle + |\downarrow_1\rangle|\uparrow_2\rangle$ are said to be **entangled**.


Experiment Two



The diagram shows a central box labeled 'source' with two arrows pointing outwards to the left and right. On the left is a box labeled '1' with a '+' sign in the top-left corner, a '-' sign in the bottom-left corner, and an upward-pointing arrow in the center. On the right is a box labeled '2' with a '-' sign in the bottom-right corner, a '+' sign in the top-right corner, and a downward-pointing arrow in the center. Below the source, two blue arrows indicate distances: one pointing left labeled '5 km' and one pointing right labeled '5 km + 1 meter'.

- **Analyzer 1** (watched by **Albert**) is placed 5 km to the left of the source.
- **Analyzer 2** (watched by **Niels**) is placed 5 km **plus one meter** to the right of the source.
- Perform **Experiment One**, exactly as before.
- How is this experiment different from the first?

Experiment Two



Albert

Niels


5 km

5 km + 1 meter

- **Albert** can tilt **Analyzer 1** any way he wants, and **Niels** can do the same with **Analyzer 2**.
- When **Analyzers 1 & 2** are tilted at different angles, they sometimes get the same answer, sometimes different answers.
- But when they compare their data, whenever the analyzers were tilted at the same angle they got **opposite** answers.
- The measurements are still 100% **anti-correlated**.

$$|\Psi_{12}\rangle = |\uparrow_1\rangle|\downarrow_2\rangle + |\downarrow_1\rangle|\uparrow_2\rangle$$

The EPR Argument



Albert

Niels

5 km

5 km + 1 meter

- **Analyzers 1 & 2** are set at the same angle and **Albert** measures the spin of **Atom 1 first**. He observes $|\uparrow_1\rangle$.
- **Albert knows** what the result of **Niels'** measurement will be **before** **Atom 2** reaches **Analyzer 2**. [And **Niels** knows he knows it.]
- ~~What if we assume locality, the Albert's measurement can't change the outcome of Niels' measurement?~~ **Niels** observes $|\downarrow_2\rangle$, and that must have been the state of **Atom 2** all along.

The EPR Argument

Albert

Niels

5 km

5 km + 1 meter

- In other words, if **Albert** can predict with 100% certainty that **Niels** will observe $|\downarrow_2\rangle$ **before** he performs the measurement, then $|\downarrow_2\rangle$ must have been the **real, definite** state of **Atom 2 at the moment the atom pair was produced**.
- Local Realism** says the atom pair was produced in the state

$$|\Psi_{12}\rangle = |\uparrow_1\rangle |\downarrow_2\rangle$$
 and the measurements revealed this **unknown reality** to us.

MAY 15, 1935

PHYSICAL REVIEW

VOLUME 47

Can Quantum-Mechanical Description of Physical Reality Be Considered Complete?

A. EINSTEIN, B. PODOLSKY AND N. ROSEN, *Institute for Advanced Study, Princeton, New Jersey*
(Received March 25, 1935)

In a complete theory there is an element corresponding to each element of reality. A sufficient condition for the reality of a physical quantity is the possibility of predicting it with certainty, without disturbing the system. In quantum mechanics in the case of two physical quantities described by non-commuting operators, the knowledge of one precludes the knowledge of the other. Then either (1) the description of reality given by the wave function in quantum mechanics is not complete or (2) these two quantities cannot have simultaneous reality. Consideration of the problem of making predictions concerning a system on the basis of measurements made on another system that had previously interacted with it leads to the result that if (1) is false then (2) is also false. One is thus led to conclude that the description of reality as given by a wave function is not complete.

Albert Einstein believed that the properties of a physical system are **objectively real** – they exist whether we measure them or not.

Einstein, Podolsky and Rosen (**EPR**) believed in the **reality** of hidden variables not described by quantum mechanics.

OCTOBER 15, 1935

PHYSICAL REVIEW

VOLUME 48

Can Quantum-Mechanical Description of Physical Reality be Considered Complete?

N. BOHR, *Institute for Theoretical Physics, University, Copenhagen*

(Received July 13, 1935)

It is shown that a certain "criterion of physical reality" formulated in a recent article with the above title by A. Einstein, B. Podolsky and N. Rosen contains an essential ambiguity when it is applied to quantum phenomena. In this connection a viewpoint termed "complementarity" is explained from which quantum-mechanical description of physical phenomena would seem to fulfill, within its scope, all rational demands of completeness.

- **The Copenhagen Interpretation** says the atom pair was produced in the **superposition** state $|\Psi_{12}\rangle = |\uparrow_1\rangle|\downarrow_2\rangle + |\downarrow_1\rangle|\uparrow_2\rangle$
- Albert's measurement of $|\uparrow_1\rangle$ **instantly collapses** $|\Psi_{12}\rangle$ into the **definite** state $|\Psi_{12}\rangle = |\uparrow_1\rangle|\downarrow_2\rangle$
- This **collapse** must be **instantaneous**, because there is no time for a signal to travel from **1** to **2**.

Albert Einstein: God does not play dice with the universe.

Niels Bohr: Who are we to tell God how to act?



Niels Bohr and Albert Einstein together at the 1930 Solvay Conference.

Philosophy or Science?

Interpretations One & Two involved *hidden variables*.

Interpretation Three said:

In general, the state of a quantum system is indeterminate until measured.

We can restate this as:

THE OUTCOME OF A QUANTUM EXPERIMENT CANNOT, ***IN GENERAL***, BE PREDICTED EXACTLY; ONLY THE PROBABILITIES OF THE VARIOUS OUTCOMES CAN BE FOUND.

Question: How comfortable are you with **Interpretation Three** (i.e. Finkelstein says Einstein is wrong and Bohr is right)?

- A. Very comfortable
- B. Getting comfortable, but still not totally convinced
- C. On the fence, I can see arguments for both sides
- D. No way, Finkelstein (and Bohr) are full of it
- E. Don't have any idea which interpretation is right

Dealing with Hidden Variables

In 1964, J. S. Bell proves *theoretically*:

No local interpretation of quantum phenomena can reproduce all of the predictions of quantum mechanics.

THE paradox of Einstein, Podolsky and Rosen [1] was advanced as an argument that quantum mechanics could not be a complete theory but should be supplemented by additional variables. These additional variables were to restore to the theory causality and locality [2]. In this note that idea will be formulated mathematically and shown to be incompatible with the statistical predictions of quantum mechanics. It is

In 1978, John Wheeler proposes ***delayed choice experiment*** to demonstrate Bell's theorem

In 1986, Alain Aspect, et al. performs set of **single photon experiments**




Bell's Theorem

There is a powerful general theorem by J. S. Bell that proves:

No local interpretation of quantum phenomena can reproduce all of the predictions of quantum mechanics.

[We can devise a realistic scheme that is non-local, but most scientists are uncomfortable with this kind of interpretation.]




Number of annual citations of "On the Einstein-Podolsky-Rosen Paradox"
J. S. Bell, *Physics* 1, 195 (1964)

Bell's Theorem

There is a more general theorem by J. S. Bell that proves:

No local interpretation of quantum phenomena can reproduce all of the predictions of quantum mechanics.




A test of Bell's Theorem performed by A. Aspect (1981)

Single Photon Experiments



“It is wrong to think that the task of physics is to find out how Nature is. Physics concerns what we can say about Nature.

– Niels Bohr



What happens when an EM wave encounters a beamsplitter?

- (A) It's reflected towards Detector A.
- (B) It's transmitted towards Detector B.
- (C) It's reflected & transmitted towards both detectors.
- (D) It alternates back and forth between the two with time.
- (E) Something else.

Animation Step-by-step Exploration quantumcurriculum.iop.org IOP Institute of Physics

Interferometer experiments with photons, particles and waves

○ Introduction
● Controls

Input Mirror 1 Beam splitter 1 Mirror 2 Detector 1 Detector 2 Coincidence counter

Detected counts

Detector 1: $N_1 = 14$
 Detector 2: $N_2 = 13$
 Coincidences: $N_c = 5$

Clear measurements

Input

Classical particles
 Electromagnetic wave
 Single photons

Main Controls

Not available for waves!
 Stop
 Insert second beam splitter

Phase shift in lower path

0 $\pi/2$ π

Display controls

Label elements
 Show theoretical intensities (Input intensity is 100%)

PHOTON SOURCE BS D-A D-B

What happens when a **single photon** encounters a beamsplitter?

(A) Either reflected towards D-A (w/ 50% probability) **or** transmitted towards D-B (w/ 50% probability) at random.
 (B) It's reflected & transmitted towards both detectors.
 (C) Science has no way of knowing.

Silent / No Discussion Please

Interpretation

Statistical: Each photon is **either** reflected **or** transmitted at the beamsplitter (but not both). The superposition state represents our ignorance of its actual state.

Quantum Wave: Each photon is **both** reflected **and** transmitted. The superposition state represents the actual state of each photon after encountering the beamsplitter.


Copenhagen: We can't describe what we can't observe. The superposition is the correct mathematical description of the possible measurement outcomes, but we can't ever know more than that.



“The result of [the detection] must be either the whole photon or nothing at all. Thus the photon must change suddenly from being partly in one beam and partly in the other to being entirely in one of the beams.”


P. A. M. Dirac, *The Principles of Quantum Mechanics* (1930, p. 8)

Single Photon Source (1986)



- Calcium atoms are excited by a two-photon absorption process ($E_K = 3.05 \text{ eV}$) + ($E_D = 2.13 \text{ eV}$).
- The excited state first decays by single photon emission ($E_1 = 2.25 \text{ eV}$).
- The lifetime of the intermediate state is $\tau \sim 5 \text{ ns}$.
- High probability the second photon ($E_2 = 2.93 \text{ eV}$) is emitted within $t = 2\tau$

EXPERIMENT ONE



EXPERIMENT ONE

If the second photon (V_2) is detected in **PMA**, then it must have been...

(A) ...reflected at BS1.
 (B) ...transmitted at BS1.
 (C) ...either reflected or transmitted at BS1.
 (D) Not enough information.

EXPERIMENT ONE

If both **PMA** and **PMB** fire within $t = 2\tau$, then the coincidence counter (**NC**) is triggered

Many photons...
 ...or classical wave-like behavior at the beamsplitter

ANTI-CORRELATION PARAMETER (α)

Want some kind of measure of how often PMA & PMB are firing simultaneously (within $t=2\tau$)

$$\alpha \equiv \frac{P_C}{P_A \cdot P_B}$$

- $P_A = \frac{N_A}{N_1}$ = probability for NA to be triggered
- $P_B = \frac{N_B}{N_1}$ = probability for NB to be triggered
- $P_C = \frac{N_C}{N_1}$ = probability for coincidence counter (Nc) to be triggered (PMA & PMB during $t=2\tau$)

ANTI-CORRELATION PARAMETER

$$\alpha \equiv \frac{P_C}{P_A \cdot P_B}$$

- If NA & NB are being triggered randomly and independently, then $\alpha = 1$ $P_C = P_A \cdot P_B$
- If NA & NB are being triggered separately (reflection **or** transmission) then $\alpha \geq 0$
 $P_C = 0$ when photons are detected by either PMA or PMB, but never both simultaneously
- If NA & NB are being triggered together (reflection **and** transmission) then $\alpha \geq 1$
 $P_C > P_A \cdot P_B$ means PMA & PMB are firing together more often than random.

